

Research article

## Absence of rapid selection for acyclovir or penciclovir resistance following suboptimal oral prodrug therapy of HSV-infected mice

Robert T Sarisky\*<sup>1</sup>, H Ron Bartus<sup>1</sup>, Shelley A Dennis<sup>1</sup>, Matthew R Quail<sup>1</sup>, Tammy T Nguyen<sup>1</sup>, Robert J Wittrock<sup>1</sup>, Wendy S Halsey<sup>2</sup>, Teresa H Bacon<sup>3</sup>, Jeffrey J Leary<sup>1</sup> and David Sutton<sup>1</sup>

Address: <sup>1</sup>Department of Host Defense, The Antimicrobial and Host Defense Center of Excellence for Drug Discovery, <sup>2</sup>Department of Discovery Genetics, GlaxoSmithKline Pharmaceuticals, Collegeville, Pennsylvania and <sup>3</sup>GlaxoSmithKline Consumer Healthcare, Weybridge, UK

E-mail: Robert T Sarisky\* - robert\_t\_sarisky@gsk.com; H Ron Bartus - henry\_ron\_bartus@gsk.com; Shelley A Dennis - shelley\_a\_dennis@gsk.com; Matthew R Quail - mquail@tektagen.com; Tammy T Nguyen - tammy\_t\_nguyen@gsk.com; Robert J Wittrock - robert\_j\_wittrock@gsk.com; Wendy S Halsey - wendy\_s\_halsey@gsk.com; Teresa H Bacon - teresa\_h\_bacon@gsk.com; Jeffrey J Leary - jjl27773@gsk.com; David Sutton - david\_sutton@gsk.com

\*Corresponding author

Published: 10 December 2001

Received: 4 October 2001

BMC Infectious Diseases 2001, 1:24

Accepted: 10 December 2001

This article is available from: <http://www.biomedcentral.com/1471-2334/1/24>

© 2001 Sarisky et al; licensee BioMed Central Ltd. Verbatim copying and redistribution of this article are permitted in any medium for any non-commercial purpose, provided this notice is preserved along with the article's original URL. For commercial use, contact [info@biomedcentral.com](mailto:info@biomedcentral.com)

### Abstract

**Background:** Acyclovir (ACV) resistant herpes simplex virus (HSV) isolates can be readily selected in animal infection models receiving suboptimal ACV treatment, however no comparative studies of the emergence of resistance following suboptimal treatment with valacyclovir (VCV) or famciclovir (FCV), the prodrugs of acyclovir and penciclovir, respectively, have been reported.

**Methods:** Mice (n = 30) were infected with HSV type 1 or 2 in the ear pinnae and administered oral prodrugs at one fifth a dose previously shown to be effective. To select and amplify drug-resistant HSV, a total of seven consecutive *in vivo* passages with suboptimal treatment were performed for each virus sample and progeny virus from each passage was characterized by the plaque reduction (PRA) and plating efficiency assays (PEA).

**Results:** No drug-resistant HSV-2 and only a single drug-resistant HSV-1 variant were identified. Virus recovered from the first three sequential passages of this HSV-1 sample was susceptible by PRA, although the proportion of resistant virus recovered gradually increased upon passage. The resistant HSV-1 phenotype was confirmed by PRA after four sequential passages in mice. Unexpectedly, this *in vivo*-selected drug-resistant HSV-1 failed to yield an infection completely refractory to treatment in subsequent passages.

**Conclusions:** Sub-optimal therapy of immunocompetent mice with either VCV or FCV did not readily select for HSV-mutants resistant to either ACV or PCV, suggesting that selection of resistance with either prodrug remains difficult using this system. Furthermore, this study suggests that the PEA may represent a useful adjunct to the PRA for monitoring alterations in the proportion of drug-resistant virus even when no change in IC<sub>50</sub> is apparent.

## Background

Resistance to acyclovir (ACV) or penciclovir (PCV) generally arises by a single base mutation in either the HSV thymidine kinase (TK) or DNA polymerase (Pol) gene [for a review, see reference [5]]. Unlike Pol, TK is not essential for virus replication in cell culture [7], although *in vivo* analyses implicate it in HSV virulence, pathogenicity and reactivation from latency [4,8,16,22]. Since ACV and PCV are dependent upon HSV TK for monophosphorylation, an event obligatory for the formation of the active triphosphate inhibitor [14], ACV<sup>r</sup> isolates often exhibit high levels of cross-resistance to PCV [2,3].

Viral populations are naturally heterogeneous; there are approximately 6 to 8 TK deficient variants per 10<sup>4</sup> plaque-forming units (PFU) of HSV-1 (laboratory strains and clinical isolates) that have never been exposed to antiviral selection [6,13,15,17]. Resistance to nucleoside inhibitors of HSV arises readily in tissue culture probably by selection of pre-existing mutants [10,13,18]. Moreover, Field *et al.*[9] demonstrated sequential HSV inoculation from mouse to mouse receiving sub-optimal ACV therapy led to the selection and amplification of resistant virus *in vivo*. After four serial passages in the immunocompetent mice, the infection became completely refractory to ACV treatment.

Acyclovir has been in part superseded by the oral prodrugs valaciclovir (VCV) and famciclovir (FCV), the latter being the oral version of PCV [20,24]. No comparative studies of the emergence of resistance following sub-optimal VCV or FCV have been reported and therefore set out to select for HSV-1 and HSV-2 isolates resistant to ACV and PCV after using prodrug selection.

## Materials and Methods

### Cell lines and virus strains

MRC-5 (pass 18–25) and Vero were grown in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal calf serum (FCS) and incubated at 37°C, 5% CO<sub>2</sub>. Vero cells were utilized for virus plaque purification and protein lysate preparations and MRC-5 cells were used for PRA and PEA. TK-negative (TK<sup>-</sup>) human osteosarcoma (143) cells were propagated in the medium described above and utilized for the <sup>3</sup>H-thymidine kinase enzymatic assay.

Virus stocks were prepared by inoculating cells at 0.01 plaque-forming units per MRC-5 cell (PFU/cell). Laboratory virus strains utilized were HSV-1 SC16 and HSV-2 SB5. SC16 was kindly provided by Dr. Sharon Safran (Gilead Sciences, CA). SB5 (ATCC; VR-2546) was plaque-purified from HSV-2 333 kindly provided by Dr. Priscilla Schaffer (University of Pennsylvania, PA) at passage 6.

### Antiviral agents

PCV and FCV were synthesized at GlaxoSmithKline. VCV was extracted from Valtrex<sup>TM</sup> tablets and purity was confirmed by HPLC.

### Animal inoculation and treatment


Female BALB/c mice (18–20 g; Charles River, Raleigh, NC) were infected intradermally in both ear pinnae with 20 µl of virus suspension containing 10<sup>5</sup> PFU of HSV as previously described [1]. The same viral challenge was used for each subsequent *in vivo* passage. VCV or FCV were administered via drinking water at either 1 mg/ml (a dose previously shown to be a highly effective [23]; and D Sutton, unpublished observations) or 0.2 mg/ml (suboptimal treatment) from the time of infection until day 4 post-infection. No significant difference in daily consumption between either of the treated groups or the placebo group (drug-free water) was observed. Daily drug intake equated to 75 mg/kg or 15 mg/kg, for the 1 mg/ml and 0.2 mg/ml doses, respectively.

To facilitate processing of blinded samples, a coding system using sample number-passage number was applied. Virus sample numbers 1 through 15 were HSV-1, and 16 through 30 were HSV-2. The ears from one mouse were pooled (passage number 1 inoculation) and used to inoculate the ear pinnae of one mouse (passage number 2). There were five replicate mice per treatment group (either suboptimally treated with 0.2 mg/ml FCV, VCV or placebo-treated) and isolates from seven serial *in vivo* passages examined. Five individual plaques were purified three times to homogeneity from the single drug-resistant virus mixture identified after *in vivo* passage, 14-P4 (sample 14, passage 4). The clonal purified isolates from this sample were designated 14-P4A-E.

The *in vivo* pathogenicity of selected virus preparations was assessed in mice by monitoring lesion severity for 14 days following inoculation of the left ear pinna with 5 × 10<sup>5</sup> PFU. Cumulative lesion scores (day 0–14) were then calculated from the individual daily scores.

### Virus isolation and *in vitro* assays

Virus stocks were prepared as described previously [17], titrated and characterized by PRA and PEA. On day 4 post-infection, mice were euthanized, and both ear pinnae removed and pooled in coded tubes, one tube per mouse. The code was not broken until all tissue culture assays had been completed. The tissue was homogenized and virus stocks were prepared in MRC-5 cells from cell-free supernatants. Following virus adsorption onto the MRC-5 monolayer, two washes with phosphate-buffered saline were performed to remove any residual antiviral drug. Cell-free virus was harvested, titrated and characterized by PRA and PEA prior to use in subsequent pas-



**Figure 1**

Experimental design for *in vivo* selection of drug-resistant HSV. This cartoon depicts the methodology used to select drug-resistant virus. Both ear pinnae of each mouse were infected with HSV. There were five replicate mice per treatment group (treated suboptimally with 0.2 mg/ml FCV or VCV, or with placebo) and isolates from seven serial *in vivo* passages examined. The resulting virus preparation from the pooled ears of an individual mouse (sample number 1, passage number 1) were used to inoculate both ear pinnae of one mouse for the subsequent serial passage (sample number 1, passage number 2).

sage experiments (Figure 1). All PRA, PEA and *in vivo* studies were carried out using virus that had undergone a single *in vitro* passage. PRAs were performed in MRC-5 cells as described previously [17]. The plating efficiency of the virus preparations generated *in vivo* was determined according to the method developed by Hall *et al.* with minor modifications [13] as reported previously [17]. Briefly, six serial 10-fold dilutions of virus were inoculated onto MRC-5 cells in the absence of antiviral drug or in the presence of either ACV or PCV at a concentration of antiviral 10 times higher than the average  $IC_{50}$  for wild type parental HSV-1 and HSV-2, respectively. The mutation frequency was calculated as follows: mutation frequency = (virus titer in the presence of drug)/(virus titer in the absence of drug). The frequency is often

expressed as a plating efficiency, or percentage of viruses that are resistant, which is calculated as percent resistant = (mutation frequency)  $\times$  100.

#### **Antibodies and biochemical assays**

Western blot analysis was performed as described previously [18]. Viral TK activity was determined by a modification of the method utilized by Coen and Schaffer [4] as described in [17]. Primers utilized for TK sequencing were described previously [18].

## Results

### Virus yield after antiviral treatment

**Table 1: Virus Titer Day 4 post-infection in mice inoculated with HSV-1 strain SC16 or HSV-2 strain SB5 and treated orally with Placebo, FCV or VCV**

Virus inoculum <sup>b</sup>	Virus Titer (log <sub>10</sub> PFU/ear)				
	Placebo	FCV		VCV	
		0.2 mg/ml <sup>c</sup>	1.0 mg/ml <sup>c</sup>	0.2 mg/ml <sup>c</sup>	1.0 mg/ml <sup>c</sup>
HSV-1 (SC16)	4.54 ± 0.04 <sup>a</sup>	2.92 ± 0.23 (2.4) <sup>d</sup>	1.0 ± 0.21 (0.03)	2.45 ± 0.15 (0.80)	0.74 ± 0.35 (0.02)
HSV-2 (SB5)	4.73 ± 0.30	2.55 ± 0.25 (0.67)	0.98 ± 0.13 (0.02)	2.06 ± 0.17 (0.20)	0.72 ± 0.44 (0.01)

<sup>a</sup>Geometric mean titer ± standard deviation of virus obtained from the ears of five mice and titrated independently. <sup>b</sup>A total of 10<sup>5</sup> PFU was inoculated into each ear pinna. <sup>c</sup>Treatment in the drinking water from the time of virus inoculation. Mice drank approximately 1.5 ml per day. No differences in intake volume or spillage between treatment groups were evident. <sup>d</sup>Percent virus yield compared to placebo-treated mice.

To evaluate efficacy of prodrug treatment in this model, VCV and FCV were provided in the drinking water at 1.0 mg/ml or at 0.2 mg/ml (sub-optimal treatment) for mice infected with either HSV-1 SC16 or HSV-2 SB5. Titers (day 4 post-infection) of approximately 4.5 log<sub>10</sub> virus per ear were recovered from placebo-treated mice, regardless of virus type (Table 1). As expected, a significant reduction in virus yield (greater than 3.5 log<sub>10</sub> decrease) was noted for mice treated with either VCV or FCV at 1.0 mg/ml following infection with either HSV-1 or HSV-2.

In the same experiment, virus yield from suboptimally-treated mice was reduced approximately 1.6–2.6 log<sub>10</sub> compared to placebo-treated mice, and these data are consistent with a previous report in which ACV was administered at a concentration of 0.25 mg/ml [10]. At 0.2 mg/ml, viral replication, albeit reduced, was still ongoing, a prerequisite for the *in vivo* selection of resistant HSV.

A reproducible difference in virus yield between the two antiviral treatment groups at the suboptimal concentration was evident, regardless of virus type. For HSV-1, the difference in titer between VCV-treated mice (2.45 log<sub>10</sub> PFU/ml) and FCV-treated (2.92 log<sub>10</sub> PFU/ml) was statistically significant ( $p = 0.049$ ). Similarly for HSV-2, VCV at 0.2 mg/ml was significantly more effective than FCV in this mouse model. Titers from VCV- or FCV-treated mice were 2.06 log<sub>10</sub> PFU/ml or 2.55 log<sub>10</sub> PFU/ml, respectively ( $p = 0.001$ ). Similar trends were noted when the antivirals were compared at 1.0 mg/ml for both HSV-1 and HSV-2 (Table 1). These observations can not

be accounted for by differences in exposure to ACV and PCV since oral administration of VCV and FCV results in almost identical blood concentration-time curves for ACV and PCV [11]. A more probable explanation is that ACV is approximately four-fold more potent than PCV in murine cells [21].

To select and amplify drug-resistant HSV in mice, a total of seven consecutive passages were performed, and progeny virus from each passage was characterized *in vitro* by PRA and PEA.

#### PRA susceptibility

IC<sub>50</sub>s were determined for each virus sample by the PRA. The parental control virus stock (which was not passaged in mice) was tested in parallel to provide a baseline for every PRA. Consistent with previous studies [2], IC<sub>50</sub>s for PCV tended to be marginally higher than for ACV, and IC<sub>50</sub>s were higher for HSV-2 SB5 than for HSV-1 SC16. The *in vitro* breakpoint for defining ACV resistance (2.0 µg/ml) and the provisionally-defined PCV resistance breakpoint, an IC<sub>50</sub> greater than 10-fold above the wild-type control virus IC<sub>50</sub> within a particular assay [17], were utilized.

#### Drug-selected HSV-1 SC16

Mean PCV susceptibilities of the HSV-1 isolates ( $n = 105$ ) from passages 1 through 7 are listed in Table 2. All isolates from mice treated with either placebo or FCV remained sensitive to PCV (IC<sub>50</sub>s ranging from 0.28–1.50 µg/ml) and to ACV (results not shown) according to the standard resistance breakpoint criteria, whereas isolates

**Table 2: Susceptibility of serial passaged HSV-1 SCI 6 isolates to PCV as measured by the PRA.**

Parental SCI 6 <sup>a</sup>	Passage No.	PCV IC <sub>50</sub> (μg/ml)		
		Placebo	FCV	VCV
0.39	1	0.36 ± 0.04 <sup>b</sup>	0.42 ± 0.07	0.42 ± 0.05
0.57	2	0.48 ± 0.16	0.76 ± 0.23	0.74 ± 0.35
0.90	3	0.33 ± 0.08	0.50 ± 0.24	0.49 ± 0.36
0.73	4	1.13 ± 0.46	1.31 ± 0.42	2.73 ± 3.11 <sup>c</sup>
0.77	5	1.15 ± 0.24	0.68 ± 0.15	0.68 ± 0.07
0.70	6	0.79 ± 0.13	0.74 ± 0.09	0.80 ± 0.12
0.70	7	0.81 ± 0.38	0.57 ± 0.27	0.66 ± 0.24

<sup>a</sup>IC<sub>50</sub> against PCV for the original parental control virus preparation (not passaged in mice). <sup>b</sup>IC<sub>50</sub> against PCV ± standard deviation for test isolates for n = 5 samples. <sup>c</sup>IC<sub>50</sub> (μg/ml) PCV for samples 26–30 ranged from 1.09–8.3.

from mice treated with VCV had PCV IC<sub>50</sub>s ranging from 0.31–8.3 μg/ml. One isolate (4th passage; 14-P4) in this treatment group was confirmed as resistant to both PCV (IC<sub>50</sub> = 8.3 μg/ml) and ACV (IC<sub>50</sub> = 9.5 μg/ml). Generally, test sample IC<sub>50</sub>s varied within two-fold of the non-passaged control virus IC<sub>50</sub>, within each particular assay (data not shown); however, isolate 14-P4 was clearly different, with an IC<sub>50</sub> greater than 11-fold over the control virus IC<sub>50</sub> for both PCV and ACV.

Interestingly, although 14-P4 was resistant, the third pass isolate for this sample was completely sensitive to both antiviral agents (ACV IC<sub>50</sub> = 0.16 μg/ml; PCV IC<sub>50</sub> = 0.53 μg/ml). Therefore, the PRA provided no early indication at pass 3 of potentially increasing resistant variants upon subsequent passage. Furthermore, the resulting virus preparation from subsequent passaging of 14-P4 reverted to a sensitive phenotype (ACV IC<sub>50</sub> = 0.74 μg/ml; PCV IC<sub>50</sub> = 0.74 μg/ml), although suboptimal treatment was still utilized.

#### Drug-selected HSV-2 SB5

Mean PCV susceptibilities of the HSV-2 isolates (n = 105) from passages 1 through 7 are listed in Table 3. No resistant HSV-2 emerged among the total of 105 HSV-2 isolates tested from the three treatment groups (PCV IC<sub>50</sub>s ranged from 0.66–1.85 μg/ml across treatment groups). These results were confirmed when the same isolates were tested for susceptibility to ACV (results not shown) and all the IC<sub>50</sub>s were sensitive over a similar range as PCV. As with HSV-1, the HSV-2 test samples were generally within two-fold of the control virus IC<sub>50</sub> (data not shown).

#### PEA profiling

All isolates were tested by the PEA, a test which may provide a more sensitive indicator of emergence of drug resistance than the PRA, since it measures the proportion of resistant isolates in the population. The parental control virus stock was tested in parallel in order to provide a baseline for every PEA.

#### Drug-selected HSV-1 SCI 6

The HSV-1 control virus preparation contained between 0.002–0.002% ACV<sup>r</sup> virus and 0.004–0.009% PCV<sup>r</sup>, which equates to an average of 6.6 ACV<sup>r</sup> and 6.0 PCV<sup>r</sup> resistant viruses per 10<sup>5</sup> PFU. Generally, the proportion of PCV<sup>r</sup> virus varied only two-fold between tests when compared to the control virus, ranging from 0.004 to 0.009% (data not shown); however, three serial samples in the VCV-treatment group, 14-P2, 14-P3 and 14-P4, clearly showed a substantial increase in the proportion of resistant virus relative to the wild type control, specifically an 11-, 300- and 8800-fold increase, respectively (14-P2, 0.002%; 14-P3, 2%; and 14-P4, 44% PCV<sup>r</sup>). Although passage 3 preparation contains a two log increase in resistant virus compared to passage 2, this level is insufficient to significantly increase the IC<sub>50</sub>. Consistent with the failure to establish an infection refractory to treatment, the proportion of resistant virus decreased to 0.08% PCV<sup>r</sup> and 0.05% ACV<sup>r</sup> by passage 5.

#### Drug-selected HSV-2 SB5

The HSV-2 control virus preparation contained between 0.24–0.40% ACV<sup>r</sup> and 0.13–0.40% PCV<sup>r</sup> virus, which equates to an average of 3.1 ACV<sup>r</sup> and PCV<sup>r</sup> viruses per 10<sup>3</sup> PFU. Generally, the proportion of resistant virus varied only three-fold between tests when compared to the control virus (data not shown), however a continual increase in the proportion of resistant virus was seen with

**Table 3: Susceptibility of serial passaged HSV-2 SB5 isolates to PCV as measured by the PRA**

Parental SB5 <sup>a</sup>	Passage No.	PCV IC <sub>50</sub> (μg/ml)		
		Placebo	FCV	VCV
0.79	1	0.76 ± 0.08 <sup>b</sup>	0.78 ± 0.05	0.87 ± 0.09
1.02	2	0.98 ± 0.08	0.82 ± 0.23	0.96 ± 0.27
1.06	3	1.10 ± 0.23	1.04 ± 0.22	0.87 ± 0.04
1.43	4	1.55 ± 0.15	1.02 ± 0.51	1.14 ± 0.58
1.11	5	1.32 ± 0.38	1.47 ± 0.48	1.14 ± 0.18
0.80	6	0.96 ± 0.18	0.81 ± 0.05	0.85 ± 0.16
0.90	7	0.81 ± 0.08	0.77 ± 0.05	0.82 ± 0.12

<sup>a</sup>IC<sub>50</sub> against PCV for the original parental control virus preparation (not passaged in mice). <sup>b</sup>IC<sub>50</sub> against PCV ± standard deviation for test isolates for n = 5 samples.

**Table 4: Biochemical Characterization of 14-P4 clonal plaque isolates.**

Clone	14-P4A	14-P4B	14-P4C	14-P4D	14-P4E	WT SC16
TK activity	0.3%	0.4%	0.5%	0.3%	0.3%	100%
PCV IC <sub>50</sub>	>25 μg/ml	>25 μg/ml	>25 μg/ml	>25 μg/ml	>25 μg/ml	0.3 μg/ml
PFA IC <sub>50</sub>	48 μg/ml	62 μg/ml	55 μg/ml	80 μg/ml	55 μg/ml	64 μg/ml

the sample 30 passages (VCV treatment group). The proportion of PCV<sup>r</sup> virus in sample 30 increased from 0.40% to 2.8% to 12.2% throughout serial passages 1 through 3 and declined to 0.2% by passage 4. All of these isolates were susceptible to PCV by the PRA (IC<sub>50</sub>s for sample 30, passage 2 through 4, were 0.79, 0.90 and 1.2 μg/ml, respectively).


#### Analysis of drug-resistant isolate

Of 210 HSV isolates from mice treated sub-optimally with FCV or VCV, or with placebo, only one HSV-1 isolate, 14-P4 was identified in the PRA as an antiviral-resistant isolate. Five single plaques from 14-P4 were isolated after three rounds of plaque purification and characterized. The clonal isolates were tested for susceptibility to using the PRA and were found to be highly resistant to ACV (ACV IC<sub>50</sub>s > 25 μg/ml) and PCV (Table 4; PCV IC<sub>50</sub>s > 25 μg/ml), although susceptible to foscarnet.

Western blot analysis showed that each clonal isolate produced a truncated TK polypeptide compared to the wild type TK protein product of approximately 43 kD (Figure 2). Moreover, all five clones were defective for TK

activity compared with HSV-1 SC16 in a <sup>3</sup>H-thymidine kinase assay (Table 4). Sequence analysis of the TK coding region from all five plaque isolates identified a G nucleotide insertion in the homopolymeric hot-spot G7 tract at nucleotide 435, a common mutation present in other ACV<sup>r</sup> HSV isolates [18,19]. This base insertion produced a frame shift in the coding sequence, resulting in a loss of the ATP / nucleoside-binding site and the creation of a premature stop codon at amino acid 228 to encode a truncated 28 kD TK polypeptide.

Lastly, the *in vivo* pathogenicities of 14-P4 and one of its clonal isolates (14-P4E) were compared with the control virus, HSV-1 SC16. Isolate 14-P4 containing 45% resistant virus caused significant clinical disease in mice, only marginally less severe than that observed with HSV-1 SC16. In marked contrast, mice inoculated with the plaque-purified isolate, 14-P4E, developed a mild infection. The cumulative disease scores for parental virus control HSV-1 SC16, 14-P4 and 14-P4E were 136, 96 and 15, respectively.



**Figure 2**

Biochemical characterization of clonal plaque isolates derived from 14-P4 in MRC-5 cells. Western blot analysis of TK protein products from five clonal isolates of virus sample 14-P4. Lanes A through E represent infected cell protein extract from isolates 14-P4A, 14-P4B, 14-P4C, 14-P4D and 14-P4E. Mock represents uninfected MRC-5 cell protein extract. Wild type TK, indicated as WT SCI16, is a 43 kD polypeptide expressed from an HSV-1 SCI16 infected cell extract.

## Discussion

HSV-1 or HSV-2 isolates were passaged serially seven times in mice inoculated in the ear pinnae and treated with suboptimal (15 mg/kg per day) concentrations of FCV or VCV to select resistant virus. Despite ongoing, albeit reduced, viral replication in the animals, selection of resistant HSV with the prodrugs proved difficult. Interestingly, the single drug-resistant isolate identified from this study reverted to a sensitive phenotype upon reinfection, even with suboptimal antiviral treatment.

Although virus recovered at passage 4 from one of five mice treated with VCV (14-P4) was clearly resistant in the PRA ( $IC_{50}$ s 8.3 and 9.5 for PCV and ACV, respectively), the preceding and the subsequent isolates were sensitive according to the criteria for defining resistance. The PEA identified a gradual increase in the proportion of resistant virus from 14-P2 (0.002%) to 14-P3 (~2%) to 14-P4 (44%). However, for the most part, the PEA data did not correlate with the eventual appearance of drug-resistant virus as defined by the PRA. Consistent with this, an earlier study showed that the proportion of drug-resistant virus in a preparation needs to be approximately 50% before resulting in a significant change in  $IC_{50}$ [9].

Biochemical characterization of the 14-P4 isolate confirmed that impaired TK activity was most likely responsible for the resistance phenotype. Sequence analysis of

one of these clonal isolates (14-P4E) revealed insertion of an extra G within the  $G_7$  homopolymeric region of the TK coding region. This mutation has also been detected in clinical isolates from patients immunocompromised [12] or with AIDS [19] as well as resistant HSV selected in tissue culture [15,18]. Interestingly, this mutation was present in ACV<sup>r</sup> clinical isolates obtained during reactivation in the absence of drug pressure, as well as in ACV<sup>r</sup> strains which evolved during ACV therapy and were followed by ACV-susceptible isolates at the next reactivation [19].

It was surprising to find that the ACV-resistant virus selected *in vivo* was not retained or amplified in subsequent passages despite continued suboptimal treatment with VCV. This result appears to be in direct contrast to that of Field [9] who found that following selection of ACV-resistant virus *in vivo*, such viruses remained refractory to treatment upon subsequent passage using suboptimal treatment with ACV. The reason for this difference is unclear. However, the failure to maintain the resistant virus on subsequent passage suggests that the pathogenicity of 14-P4 may have been only partially attenuated relative to the sensitive virus and indeed the pathogenicity of the clonal virus isolate, 14-P4E, was severely attenuated. Further studies on the 14-P4 clonal isolates and the clinical significance of their attenuation should be explored.

## Conclusions

Sub-optimal therapy of immunocompetent mice with either VCV or FCV did not readily select for HSV-mutants resistant to either ACV or PCV, suggesting that selection of resistance with either prodrug remains difficult using this system. All the virus isolates from each serial passage were examined by both PRA and PEA. The former is widely accepted as the gold standard for susceptibility testing of HSV, and for ACV, a correlation between PRA *in vitro*  $IC_{50}$  and clinical outcome has been established. In spite of this, the PEA appears to have potential as an adjunct to the PRA by providing earlier detection of the emergence of resistant HSV. However, it is important to note that although an increase in the number resistant variants within isolates was readily detected by the PEA, clinically-resistant virus did not develop in the immunocompetent mice. Therefore, the presence of resistant virus within an isolate, even at levels substantially above background, does not in itself indicate that increased resistance will follow, nor does it necessarily predict clinical resistance. It would be of interest to apply the PEA to an extensive series of clinical isolates, particularly isolates obtained from severely immunocompromised patients, in order to evaluate the proportions of resistant virus that occur within different clinical isolates.

## Competing Interests

None declared.

## Acknowledgements

We thank S. Albelda, S. Safrin and P. Schaffer for generous gifts of reagents; P. Patel and N. Simon for technical assistance; B. Gagnon for statistical analyses; R. Boon for financial support of the Consumer Healthcare division; SB Laboratory Animal Science for expert handling and care of animals in accordance with the guidelines of the SB Animal Care and Use Committee; and K. Esser for support.

## References

- Boyd MR, Bacon TH, Sutton D: **Antiherpesvirus activity of 9-(4-hydroxy-3-hydroxymethylbut-1-yl) guanine (BRL 39123) in animals.** *Antimicrob Agents Chemother* 1988, **32**:358-363
- Boyd MR, Bacon TH, Sutton D, Cole M: **Antiherpesvirus activity of 9-(4-hydroxy-3-hydroxymethylbut-1-yl)guanine (BRL 39123) in cell culture.** *Antimicrob Agents Chemother* 1987, **31**:1238-1242
- Boyd MR, Safrin S, Kern ER: **Penciclovir – a review of its spectrum of activity, selectivity, and cross-resistance pattern.** *Antiviral Chem Chemother* 1993, **4**:3-11
- Coen DM, Kosz-Vnenchak M, Jacobson JG, Leib DA, Bogard CL, Schaffer PA, Tyler KL, Knipe DM: **Thymidine kinase-negative herpes simplex virus mutants establish latency in mouse trigeminal ganglia but do not reactivate.** *Proc Natl Acad Sci U.S.A* 1989, **86**:4736-4740
- Collins P, Darby G: **Laboratory studies of herpes simplex virus strains resistant to acyclovir.** *Rev. Med. Virol* 1991, **1**:19-28
- Dasgupta UB, Summers WC: **Ultraviolet reactivation of herpes simplex virus is mutagenic and inducible in mammalian cells.** *Proc Natl Acad Sci U.S.A* 1978, **75**:2378-2381
- Dubbs DR, Kit S: **Mutant strains of herpes simplex deficient in thymidine kinase-inducing activity.** *Virology* 1964, **22**:493-502
- Efstathiou S, Kemp S, Darby G, Minson AC: **The role of herpes simplex virus type I thymidine kinase in pathogenesis.** *J Gen Virol* 1989, **70**:869-879
- Field HJ: **Development of clinical resistance to acyclovir in herpes simplex virus-infected mice receiving oral therapy.** *Antimicrob Agents Chemother* 1982, **21**:744-752
- Field HJ, Darby G, Wildy P: **Isolation and characterization of acyclovir-resistant mutants of herpes simplex virus.** *J Gen Virol* 1980, **49**:115-124
- Field HJ, Tewari D, Sutton D, Thackray AM: **Comparison of efficacies of famciclovir and valaciclovir against herpes simplex virus type I in a murine immunosuppression model.** *Antimicrob Agents Chemother* 1995, **39**:1114-1119
- Gaudreau A, Hill E, Balfour HH, Erice A, Boivin G: **Phenotypic and genotypic characterization of acyclovir-resistant herpes simplex viruses from immunocompromised patients.** *J Infect Dis* 1998, **178**:297-303
- Hall JD, Coen DM, Fisher BL, Weisslitz M, Randall S, Almy RE, Gelep PT, Schaffer PA: **Generation of genetic diversity in herpes simplex virus: an antimutator phenotype maps to the DNA polymerase locus.** *Virology* 1984, **132**:26-37
- Hill EL, Hunter GA, Ellis MN: **In vitro and in vivo characterization of herpes simplex virus clinical isolates recovered from patients infected with human immunodeficiency virus.** *Antimicrob Agents Chemother* 1991, **35**:2322-2328
- Hwang CC, Chen HH: **An altered spectrum of herpes simplex virus mutations mediated by an antimutator DNA polymerase.** *Gene* 1995, **152**:191-193
- Jacobson JG, Ruffner KL, Kosz-Vnenchak M, Hwang CBC, Knipe DM, Coen DM: **Herpes simplex virus thymidine kinase and specific stages of latency in murine trigeminal ganglia.** *J Virol* 1993, **67**:6903-6908
- Sarisky RT, Nguyen TT, Duffy KE, Wittrock RJ, Leary JJ: **Difference in incidence of spontaneous mutations distinct between herpes simplex virus types I and 2.** *Antimicrob Agents Chemother* 2000, **44**:1524-1529
- Sarisky RT, Quail MR, dark PE, Nguyen TT, Halsey WS, Wittrock RJ, Bartus JO, Van Horn MM, Sathe GS, Van Horn S, Kelly MD, Bacon TH, Leary JJ: **Characterization of herpes simplex viruses selected in culture for resistance to penciclovir or acyclovir.** *J Virol* 2001, **75**:1761-1769
- Sasadeusz JJ, Tufaro F, Safrin S, Schubert K, Hubinette MM, Cheung PK, Sacks S: **Homopolymer mutational hot spots mediate herpes simplex virus resistance to acyclovir.** *J Virol* 1997, **71**:3872-3878
- Soullawton J, Seaber E, On N, Wootton R, Rolan P, Posner J: **Absolute bioavailability and metabolic disposition of valaciclovir, the l-valyl ester of acyclovir, following oral administration to humans.** *Antimicrob Agents Chemother* 1995, **39**:2759-2764
- Sutton D, Boyd MR: **Comparative activity of penciclovir and acyclovir in mice infected intraperitoneally with herpes simplex virus type I SC16.** *Antimicrob Agents Chemother* 1993, **37**:642-645
- Tenser RB, Edris WA: **Trigeminal ganglion infection by thymidine kinase-negative mutants of herpes simplex virus after in vivo complementation.** *J Virol* 1987, **61**:2171-2174
- Thackray AM, Field HJ: **Differential effects of famciclovir and valaciclovir on the pathogenesis of herpes simplex virus in a murine infection model including reactivation from latency.** *J Infect Dis* 1996, **173**:291-299
- Vere HR, Sutton D, Boyd MR, Harnden MR, Jarvest RL: **Selection of an oral prodrug (BRL 42810; famciclovir) for the antiherpesvirus agent BRL 39123 [9-(4-hydroxy-3-hydroxymethylbut-1-yl)guanine; penciclovir].** *Antimicrob Agents Chemother* 1989, **33**:1765-1773

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMedcentral will be the most significant development for disseminating the results of biomedical research in our lifetime."

Paul Nurse, Director-General, Imperial Cancer Research Fund

Publish with **BMC** and your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours - you keep the copyright

Submit your manuscript here:

<http://www.biomedcentral.com/manuscript/>



[editorial@biomedcentral.com](mailto:editorial@biomedcentral.com)