

### **EPOSTER PRESENTATION**

**Open Access** 

# Assessment of oxidative stress parameters in HIV infection

Shreewardhan Rajopadhye<sup>1,2\*</sup>, A Rosalind Marita<sup>1</sup>, MA Ansari<sup>3</sup>, Abhay Chowdhary<sup>1</sup>, Sucheta Dandekar<sup>2</sup>

From 2nd International Science Symposium on HIV and Infectious Diseases (HIV SCIENCE 2014) Chennai, India. 30 January - 1 February 2014

#### **Background**

Both viral and host factors are responsible for oxidative stress in HIV disease, which in turn activates the replication of HIV provirus by various pathways. Oxidizing stress is a pathologic phenomenon resulting from imbalance between the system producing active oxygen species and those defending the organism. The present study was aimed to assess oxidative stress markers in HIV patients.

#### **Methods**

The study included 30 HIV sero-positive patients, 30 healthy volunteers served as controls. Patients were categorized on the basis of their absolute CD4 counts into 3 groups - Group-1 (>500 CD4 cells/mm³), Group-2 (200–499 CD4 cells/mm³), and Group-3 (<200 CD4 cells/mm³). Lipid peroxidation was estimated using serum malondialdehyde as a marker, serum nitric oxide levels were assessed by Griess reagent method, serum reduced GSH by Beutler *et al*, serum C reactive protein, serum AOPP by Witko Savark method and serum proteins by Bradford method. Statistical analysis was done using the Student's t test and one-way ANOVA.

#### **Results**

Significant decrease (p<0.004) in GSH levels and significant increase (p<0.0008) in NO levels was observed in HIV infected group when compared to controls. However, no significant changes were found in levels of AOPP, MDA, and CRP in the study groups. Significant increase (p<0.0001) in MDA levels in group 3and in GSH levels (p<0.0395) in all 3 groups was seen as compared to controls.

Full list of author information is available at the end of the article

#### **Conclusion**

The findings indicate that considerable amount of oxidative stress are induced and changes in NO and GSH levels may contribute to the immunopathophysiology during HIV infection.

#### Authors' details

<sup>1</sup>Department of Biochemistry, Haffkine Institute, Mumbai-400012, Maharashtra, India. <sup>2</sup>Department of Biochemistry, Seth G.S.Medical College & KEM Hospital, Parel, Mumbai-400012, Maharashtra, India. <sup>3</sup>Department of Pathology, Seth G.S.Medical College & KEM Hospital, Parel, Mumbai-400012, Maharashtra, India.

Published: 27 May 2014

doi:10.1186/1471-2334-14-\$3-F28

Cite this article as: Rajopadhye et al.: Assessment of oxidative stress parameters in HIV infection. BMC Infectious Diseases 2014 14(Suppl 3):E28

## Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at www.biomedcentral.com/submit





<sup>\*</sup> Correspondence: shrajopadhye@gmail.com

<sup>&</sup>lt;sup>1</sup>Department of Biochemistry, Haffkine Institute, Mumbai-400012, Maharashtra. India